



## AP<sup>®</sup> Physics B 2001 Sample Student Responses

**The materials included in these files are intended for non-commercial use by AP teachers for course and exam preparation; permission for any other use must be sought from the Advanced Placement Program. Teachers may reproduce them, in whole or in part, in limited quantities, for face-to-face teaching purposes but may not mass distribute the materials, electronically or otherwise. These materials and any copies made of them may not be resold, and the copyright notices must be retained as they appear here. This permission does not apply to any third-party copyrights contained herein.**

These materials were produced by Educational Testing Service (ETS), which develops and administers the examinations of the Advanced Placement Program for the College Board. The College Board and Educational Testing Service (ETS) are dedicated to the principle of equal opportunity, and their programs, services, and employment policies are guided by that principle.

The College Board is a national nonprofit membership association dedicated to preparing, inspiring, and connecting students to college and opportunity. Founded in 1900, the association is composed of more than 3,900 schools, colleges, universities, and other educational organizations. Each year, the College Board serves over three million students and their parents, 22,000 high schools, and 3,500 colleges, through major programs and services in college admission, guidance, assessment, financial aid, enrollment, and teaching and learning. Among its best-known programs are the SAT<sup>®</sup>, the PSAT/NMSQT<sup>™</sup>, the Advanced Placement Program<sup>®</sup> (AP<sup>®</sup>), and Pacesetter<sup>®</sup>. The College Board is committed to the principles of equity and excellence, and that commitment is embodied in all of its programs, services, activities, and concerns.

Copyright © 2001 by College Entrance Examination Board. All rights reserved. College Board, Advanced Placement Program, AP, and the acorn logo are registered trademarks of the College Entrance Examination Board.


# PHYSICS B

## SECTION II

Time—90 minutes

7 Questions



**Directions:** Answer all seven questions, which are weighted according to the points indicated. The suggested time is about 15 minutes for answering each of questions 1-4, and about 10 minutes for answering each of questions 5-7. The parts within a question may not have equal weight. Show all your work in this booklet in the spaces provided after each part, NOT in the green insert.



1. (15 points) "bonus"


A ball of mass  $M$  is attached to a string of length  $R$  and negligible mass. The ball moves clockwise in a vertical circle, as shown above. When the ball is at point  $P$ , the string is horizontal. Point  $Q$  is at the bottom of the circle and point  $Z$  is at the top of the circle. Air resistance is negligible. Express all algebraic answers in terms of the given quantities and fundamental constants.

- (a) On the figures below, draw and label all the forces exerted on the ball when it is at points  $P$  and  $Q$ , respectively.



GO ON TO THE NEXT PAGE.

- (b) Derive an expression for  $v_{\min}$ , the minimum speed the ball can have at point Z without leaving the circular path.



$$(m)F_c = (m)\frac{v^2}{r}$$

$$F_c = \frac{mv^2}{r}$$

$$F_c = W = Mg$$

$$(M)(g) = \frac{Mv^2}{r}$$

$$v = \sqrt{gr}$$

- (c) The maximum tension the string can have without breaking is  $T_{\max}$ . Derive an expression for  $v_{\max}$ , the maximum speed the ball can have at point Q without breaking the string.

$$T_{\max} - Mg = \frac{Mv^2}{r}$$

$$\frac{r(T_{\max} - Mg)}{M} = v^2$$

$$v_{\max} = \sqrt{\frac{r(T_{\max} - Mg)}{M}}$$

- (d) Suppose that the string breaks at the instant the ball is at point P. Describe the motion of the ball immediately after the string breaks.




it goes straight upward, accelerating ~~at~~  
 $\text{at } -9.8 \text{ m/s}^2$

GO ON TO THE NEXT PAGE.

**PHYSICS B**  
**SECTION II**  
Time—90 minutes  
7 Questions



**Directions:** Answer all seven questions, which are weighted according to the points indicated. The suggested time is about 15 minutes for answering each of questions 1-4, and about 10 minutes for answering each of questions 5-7. The parts within a question may not have equal weight. Show all your work in this booklet in the spaces provided after each part, NOT in the green insert.



1. (15 points)


A ball of mass  $M$  is attached to a string of length  $R$  and negligible mass. The ball moves clockwise in a vertical circle, as shown above. When the ball is at point  $P$ , the string is horizontal. Point  $Q$  is at the bottom of the circle and point  $Z$  is at the top of the circle. Air resistance is negligible. Express all algebraic answers in terms of the given quantities and fundamental constants.

- (a) On the figures below, draw and label all the forces exerted on the ball when it is at points  $P$  and  $Q$ , respectively.



GO ON TO THE NEXT PAGE.

- (b) Derive an expression for  $v_{\min}$ , the minimum speed the ball can have at point Z without leaving the circular path.



$$F = ma$$

$$F = \frac{mv^2}{r}$$

$$a = \frac{v^2}{r}$$

$$T + Mg = \frac{mv^2}{r}$$

$$mv^2 = (T + Mg)r$$

$$v^2 = \frac{rT + rMg}{m}$$

$$v_{\min} = \sqrt{\frac{rT + rMg}{m}}$$

- (c) The maximum tension the string can have without breaking is  $T_{\max}$ . Derive an expression for  $v_{\max}$ , the maximum speed the ball can have at point Q without breaking the string.

$$F = ma$$

$$a = \frac{v^2}{r}$$

$$F = \frac{mv^2}{r}$$

$$T_{\max} - Mg = \frac{Mv^2}{r}$$

$$rT_{\max} - rMg = Mv^2$$

$$v^2 = \frac{rT_{\max} - rMg}{M}$$

$$v_{\max} = \sqrt{\frac{rT_{\max} - rMg}{M}}$$

- (d) Suppose that the string breaks at the instant the ball is at point P. Describe the motion of the ball immediately after the string breaks.

The ball will travel straight up with an initial velocity that is between  $v_{\min}$  and  $v_{\max}$ , and its acceleration will be  $g$  ( $9.8 \frac{m}{s^2}$ ) down, so eventually the ball will fall straight back down to Earth.

GO ON TO THE NEXT PAGE.