Harold's Physics Cheat Sheet

5 April 2023

	Mechanics: Linear Translation	Mechanics: Angular / Rotational Motion	Electricity / Magnetism	Fluid Mechanics / Thermodynamics	Atomic and Nuclear / Waves and Optics
Kinematics					
Position (m) (rad)	*Horizontal / 1-D:* $\boldsymbol{x} = \boldsymbol{x}_0 + \boldsymbol{v}_{x0}t + \frac{1}{2}\boldsymbol{a}t^2$ *Vertical:* $\boldsymbol{y} = \boldsymbol{y}_0 + \boldsymbol{v}_{y0}t - \frac{1}{2}\boldsymbol{g}t^2$ $\boldsymbol{x} = \boldsymbol{x}_0 + \boldsymbol{v}t$ $\boldsymbol{x} = \int \boldsymbol{v}\,dt$ $s = r\theta$	$\boldsymbol{\theta} = \boldsymbol{\theta}_0 + \boldsymbol{\omega}_0 t + \frac{1}{2}\boldsymbol{\alpha}t^2$ $\boldsymbol{\theta} = \boldsymbol{\theta}_0 + \boldsymbol{\omega}t$ $\boldsymbol{\theta} = \int \boldsymbol{\omega}\,dt$	$10^{-100} = googolth$ $10^{-24} = yocto-$ $10^{-21} = zepto-$ $10^{-18} = atto-$ $10^{-15} = femto-$ $10^{-12} = pico-$ $10^{-9} = nano-$ $10^{-6} = micro-$ $10^{-3} = milli-$ $10^{-2} = centi-$ $10^{-1} = deci-$ $10^{0} = 1$ $10^{1} = deca-$ $10^{2} = hecto-$ $10^{3} = kilo-$ $10^{6} = mega-$ $10^{9} = giga-$ $10^{12} = tera-$ $10^{15} = peta-$ $10^{18} = exa-$ $10^{21} = zetta-$ $10^{24} = yotta-$ $10^{100} = googol$ $10^{1000} = googolplex$	*Fluid Mechanics:* $P_1 + \rho g y_1 + \frac{1}{2}\rho v_1^2 = P_2 + \rho g y_2 + \frac{1}{2}\rho v_2^2$ *(Conservation of Mass)* $\rho = \frac{m}{V}$ $\Delta\ell = \alpha \ell_0 \Delta T$	*Waves:* $f(x,t) = A\sin\left(2\pi\left(ft - \frac{x}{\lambda}\right) + \phi\right) + k$ *Optics:* $f = \frac{v}{\lambda}$ $\frac{1}{d_o} + \frac{1}{d_i} = \frac{1}{f}$ ***Refraction:*** *(bend)* $n = \frac{c}{v}$ *Snell's Law:* $n_1 \sin\theta_1 = n_2 \sin\theta_2$ $\frac{n_1}{n_2} = \frac{v_2}{v_1}$ ***Diffraction:*** *(spread out)* $\Delta L = d\sin\theta$ $m\lambda = d\sin\theta$
	$\boldsymbol{x}(t) = A\cos(\boldsymbol{\omega}t + \phi)$ $\boldsymbol{x}(t) = A\cos(2\pi f t + \phi)$				

Copyright © 2011-2023 by Harold Toomey, Wyzant Tutor

	Mechanics: Linear	Mechanics: Angular	Electricity / Magnetism	Fluid Mechanics / Thermodynamics	Atomic and Nuclear / Waves and Optics
Velocity (m/s) Angular Velocity / Angular Frequency (rad/s)	$\boldsymbol{v} = \frac{d}{t} = \frac{\Delta \boldsymbol{x}}{\Delta t} \rightarrow \frac{d\boldsymbol{x}}{dt}$ $\boldsymbol{v} = \boldsymbol{v}_0 + \boldsymbol{a}t$ $\boldsymbol{v}^2 = \boldsymbol{v}_0^2 + 2\boldsymbol{a}(x - x_0)$ $\bar{\boldsymbol{v}} = \frac{\boldsymbol{v}_0 + \boldsymbol{v}}{2}$ $\boldsymbol{v} = \int \boldsymbol{a}\, dt$ $f = \frac{1}{T}$	$\boldsymbol{\omega} = \frac{\theta}{t} = \frac{\Delta \theta}{\Delta t} \rightarrow \frac{d\boldsymbol{\theta}}{dt}$ $\boldsymbol{\omega} = \boldsymbol{\omega}_0 + \boldsymbol{\alpha}t$ $\boldsymbol{\omega}^2 = \boldsymbol{\omega}_0^2 + 2\boldsymbol{\alpha}(\theta - \theta_0)$ $\bar{\boldsymbol{\omega}} = \frac{\boldsymbol{\omega}_0 + \boldsymbol{\omega}}{2}$ $\boldsymbol{\omega} = \int \boldsymbol{\alpha}\, dt$ $\boldsymbol{\omega} = \frac{2\pi}{T} = 2\pi f$ $\boldsymbol{\omega} = \sqrt{\frac{k}{m}}\ (spring)$ $\boldsymbol{\omega} = \sqrt{\frac{\boldsymbol{g}}{\ell}}\ (pendulum)$	*Speed of Light:* $c \approx 3.00 \times 10^8\ \frac{m}{s}$	*<u>Fluid Mechanics</u>:* $A_1 v_1 = A_2 v_2$ $v_{rms} = \sqrt{\frac{3RT}{M}}$ $v_{rms} = \sqrt{\frac{3k_B T}{\mu}}$	*<u>Waves and Optics</u>:* $v = f\lambda$ ***Reflection**:* *(throw back)* *Critical angle:* $\sin\theta_c = \frac{n_1}{n_2}$ *Maxima for a thin film:* $2d = \frac{\lambda}{2n}, 3\frac{\lambda}{2n}, 5\frac{\lambda}{2n} \ldots$
	$\boldsymbol{v} = \boldsymbol{\omega} r$ $\boldsymbol{v} = \boldsymbol{\omega} \times \boldsymbol{r}$ $\boldsymbol{v}(t) = -A\boldsymbol{\omega}\sin(\omega t + \phi)$ $\boldsymbol{v}(t) = -A(2\pi f)\sin(2\pi f t + \phi)$				

Copyright © 2011-2023 by Harold Toomey, Wyzant Tutor

	Mechanics: Linear	Mechanics: Angular	Electricity / Magnetism	Fluid Mechanics / Thermodynamics	Atomic and Nuclear / Waves and Optics
Acceleration **(m/s^2)** (rad/s^2)	*Linear:* $\boldsymbol{a} = \frac{\boldsymbol{v}}{t} = \frac{\Delta \boldsymbol{v}}{\Delta t} \rightarrow \frac{d\boldsymbol{v}}{dt}$ $\boldsymbol{a} = \frac{\sum \boldsymbol{F}}{m} = \frac{\boldsymbol{F}_{net}}{m}$ *Tangential (linear):* $\boldsymbol{a} = \boldsymbol{a}_t = \boldsymbol{\alpha}\, r$ $Gravity\ (\boldsymbol{g})$	*Angular:* $\boldsymbol{\alpha} = \frac{\boldsymbol{\omega}}{t} = \frac{\Delta \boldsymbol{\omega}}{\Delta t} \rightarrow \frac{d\boldsymbol{\omega}}{dt}$ $\boldsymbol{\alpha} = \frac{\sum \boldsymbol{\tau}}{I} = \frac{\boldsymbol{\tau}_{net}}{I}$ *Centripetal (center):* $\boldsymbol{a}_c = \frac{\boldsymbol{v}^2}{r} = \boldsymbol{\omega}^2 r$	**Constants:** *Gravitational Constant* $G = 6.67430 \times 10^{-11}\ \frac{Nm^2}{kg}$ *Gravity Acceleration (Earth)* $g = 9.80665\ \frac{m}{s^2} = 32.1740\ \frac{ft}{s^2}$ *Speed of Light in Vacuum* $c = 2.99792458 \times 10^8\ \frac{m}{s}$ *Electron-Volt* $1eV = 1.602176634 \times 10^{-19}\ J$ *Charge of an Electron* $e = -1.602176634 \times 10^{-19}\ C$ *Mass of an Electron* $m_e = 9.1093837015(28) \times 10^{-31}\ kg$ *Mass of a Proton* $m_p = 1.67262192369(51) \times 10^{-27}\ kg$ *Mass of a Neutron* $m_n = 1.67492749804(95) \times 10^{-27}\ kg$ *Electric Permittivity* $\varepsilon_0 = \frac{1}{\mu_0 c^2} = 8.8541878128 \times 10^{-12}\ \frac{C^2}{Nm^2}$ *Magnetic Permeability* $\mu_0 = 4\pi \times 10^{-7}\ \frac{Tm}{A}$ *Avogadro's Number* $N_A = 6.02214076 \times 10^{23}\ \frac{1}{mol}$ *Boltzmann Constant* $k_B = 1.380649 \times 10^{-23}\ \frac{J}{K}$ *Coulomb Constant* $k_e = \frac{1}{4\pi\varepsilon_0} = 8.9875517923 \times 10^9\ \frac{Nm^2}{C^2}$ *Faraday Constant* $F = 9.64853321233100184 \times 10^4\ \frac{C}{mol}$ *Planck's Constant* $h = 6.6267015 \times 10^{-34}\ Js$ *Avogadro's Number* $N_A = 6.02214076 \times 10^{23}\ /\ mole$ *Pi* $\pi = 3.14159265358979323846\ldots$		
	 *Net:* $\boldsymbol{a}_{net}^2 = \boldsymbol{a}_t^2 + \boldsymbol{a}_c^2$				
	$\boldsymbol{a}(t) = -A\boldsymbol{\omega}^2 \cos(\omega t + \phi)$ $\boldsymbol{a}(t) = -A(2\pi f)^2 \cos(2\pi f t + \phi)$				
Jerk (Jolt) **(m/s^3)** (rad/s^3)	$\vec{\boldsymbol{j}} = \frac{\boldsymbol{a}}{t} = \frac{\Delta \boldsymbol{a}}{\Delta t} \rightarrow \frac{d\boldsymbol{a}}{dt}$	$\boldsymbol{\zeta} = \frac{\boldsymbol{\alpha}}{t} = \frac{\Delta \boldsymbol{\alpha}}{\Delta t} \rightarrow \frac{d\boldsymbol{\alpha}}{dt}$			

Copyright © 2011-2023 by Harold Toomey, Wyzant Tutor

	Mechanics: Linear	Mechanics: Angular	Electricity / Magnetism	Fluid Mechanics / Thermodynamics	Atomic and Nuclear / Waves and Optics
Dynamics					
Mass (kg) / Moment of Inertia $(kg \bullet m^2)$	m = actual mass I = effective mass	$I = \sum mr^2$ $I = \int r^2\, dm$ $I = \int \boldsymbol{r} \bullet \boldsymbol{r}\, dm$	$m_p \approx m_n$	*NA*	***Magnification***: $\lvert M \rvert = \left\lvert \frac{d_i}{d_o} \right\rvert = \left\lvert \frac{h_i}{h_o} \right\rvert$ $M_{Telescope} = -\frac{f_{objective\ lens}}{f_{eyepiece}}$
Momentum (kg•m/s) $(kg \bullet m^2/s)$	$\boldsymbol{p} = m\boldsymbol{v}$ $\Delta\boldsymbol{p} = m\Delta\boldsymbol{v}$ *Conservation of Linear Momentum:* $\boldsymbol{p}_i = \boldsymbol{p}_f$ $\sum m\boldsymbol{v}_i = \sum m\boldsymbol{v}_f$ $\boldsymbol{p} = \boldsymbol{\omega} \times \boldsymbol{m}$ $\Delta\boldsymbol{p} = \boldsymbol{J} = \int_{t_1}^{t_2} \boldsymbol{F}\, dt$ *Elastic Collision = bounce off* *Inelastic Collision = stick together*	$\boldsymbol{L} = I\boldsymbol{\omega}$ $\boldsymbol{L} = \boldsymbol{r}m\boldsymbol{v}$ *Conservation of Angular Momentum:* $\boldsymbol{L}_i = \boldsymbol{L}_f$ $\sum I\boldsymbol{\omega}_i = \sum I\boldsymbol{\omega}_f$ $\boldsymbol{L} = \boldsymbol{r} \times \boldsymbol{p}$ $\boldsymbol{L} = m\boldsymbol{r} \times \boldsymbol{v}$ $L = \int \boldsymbol{r} \times \boldsymbol{v}\, dm$ $\Delta\boldsymbol{L} = \int \boldsymbol{\tau}\, dt$	*NA*	*<u>Fluid Mechanics</u>:* $\nabla\boldsymbol{p} = \rho\boldsymbol{g}$	*<u>Atomic and Nuclear</u>:* $\lambda = \frac{h}{p}$



Copyright © 2011-2023 by Harold Toomey, Wyzant Tutor

	Mechanics: Linear	Mechanics: Angular	Electricity / Magnetism	Fluid Mechanics / Thermodynamics	Atomic and Nuclear / Waves and Optics
Force (N = kg•m/s^2) / Torque (J = N•m)	$\boldsymbol{F} = m\boldsymbol{a}$ $\boldsymbol{F}_g = m\boldsymbol{g}$ $\sum \boldsymbol{F} = \boldsymbol{F}_{net} = m\boldsymbol{a}$ $\boldsymbol{F} = \frac{\boldsymbol{p}}{t} = \frac{\Delta p}{\Delta t}$ $\boldsymbol{F} = \frac{d\boldsymbol{p}}{dt} = \frac{d}{dt}(m\boldsymbol{v})$ $\boldsymbol{F}_f \leq \mu \boldsymbol{N}$ *Hooke's Law:* $\boldsymbol{F}_s = -k\boldsymbol{\Delta x}$ $F_G = -\frac{Gm_1m_2}{r^2}$ $\boldsymbol{F}_G = -\frac{Gm_1m_2}{r^2}\hat{\boldsymbol{r}}$	$\tau = rF\sin\theta$ $\sum \boldsymbol{\tau} = \boldsymbol{\tau}_{net} = I\boldsymbol{\alpha}$ $\tau = r\boldsymbol{F} = I\boldsymbol{\alpha}$ $\boldsymbol{\tau} = \boldsymbol{r} \times \boldsymbol{F}$ $\boldsymbol{\tau} = \frac{d\boldsymbol{L}}{dt} = I\frac{d\boldsymbol{\omega}}{dt}$ $\boldsymbol{F} = m\boldsymbol{a_c}$ $\boldsymbol{F} = \frac{m\boldsymbol{v}^2}{r}$ $\boldsymbol{F} = mr\boldsymbol{\omega}^2$	*Electricity:* *Coulomb's Law:* $\boldsymbol{F} = k\frac{q_1q_2}{\boldsymbol{r}^2}$ $\boldsymbol{F} = \frac{1}{4\pi\epsilon_0}\frac{q_1q_2}{\boldsymbol{r}^2}$ $\boldsymbol{F} = \boldsymbol{E}\,q$ *Magnetism:* $F_B = qvB\sin\theta$ $F_B = BI\ell\sin\theta$ $\boldsymbol{F}_B = q\boldsymbol{v} \times \boldsymbol{B}$ $\boldsymbol{F}_B = I\boldsymbol{\ell} \times \boldsymbol{B}$ $\boldsymbol{F} = \int I\,d\boldsymbol{\ell} \times \boldsymbol{B}$	*Fluid Mechanics:* $F = PA$ $P = \frac{F}{A} = \rho hg$ $F_{buoy} = \rho Vg$ $PV = nRT = Nk_BT$ $R = 8.31\frac{J}{(mol \bullet K)}$ $k_B = 1.38 \times 10^{-23}\frac{J}{K}$	*NA*
Impulse (N•s) (N•m•s)	$\boldsymbol{J} = \boldsymbol{F}\,\Delta t = \Delta\boldsymbol{p} = m\,\Delta\boldsymbol{v}$ $\boldsymbol{J} = \Delta\boldsymbol{p} = \int \boldsymbol{F}\,dt$	$\boldsymbol{H} = \boldsymbol{\tau}\,\Delta t = \Delta\boldsymbol{L} = I\,\Delta\boldsymbol{\omega}$ $\boldsymbol{H} = \Delta\boldsymbol{L} = \int \boldsymbol{\tau}\,dt$	*NA*	*NA*	*NA*
Yank (N/s^2) / Rotatum (J/s)	$\boldsymbol{Y} = m\boldsymbol{J}$ $\boldsymbol{Y} = \frac{\boldsymbol{F}}{t} = \frac{\Delta F}{\Delta t} \rightarrow \frac{d\boldsymbol{F}}{dt}$	$\boldsymbol{P} = I\,\boldsymbol{\zeta}$ $\boldsymbol{P} = \boldsymbol{r} \times \boldsymbol{Y}$ $\boldsymbol{P} = \frac{\boldsymbol{\tau}}{t} = \frac{\Delta\tau}{\Delta t} \rightarrow \frac{d\boldsymbol{\tau}}{dt} = I\boldsymbol{\zeta}$	*NA*	*NA*	*NA*

Copyright © 2011-2023 by Harold Toomey, Wyzant Tutor

	Mechanics: Linear	Mechanics: Angular	Electricity / Magnetism	Fluid Mechanics / Thermodynamics	Atomic and Nuclear / Waves and Optics
Energy					
Work (J = N•m)	$W = Fd$ $W = F\,\Delta x \cos\theta$ $W = \int \boldsymbol{F} \bullet d\boldsymbol{r}$	$W = \boldsymbol{\tau}\,\Delta\theta(\theta - \theta_0)$ $W = \tau\,\Delta\theta \cos\theta$ $W = \int \boldsymbol{\tau} \bullet d\boldsymbol{\theta}$	$\boldsymbol{W} = \boldsymbol{Q}V$	*Thermodynamics:* $W = -P\,\Delta V$ $e = \left\lvert \frac{W}{Q_H} \right\rvert$ $e_c = \frac{T_H - T_C}{T_H}$	*NA*
Kinetic Energy (J)	*Translational:* $K = KE = \frac{1}{2}m\boldsymbol{v}^2$	*Rotational:* $K = KE = \frac{1}{2}I\boldsymbol{\omega}^2$	$1\ eV$ $= 1.60 \times 10^{-19}\ J$	*Fluid Mechanics:* *Bernoulli's Equation:* $P_1 + \rho g y_1 + \frac{1}{2}\rho {v_1}^2$ $= P_2 + \rho g y_2 + \frac{1}{2}\rho {v_2}^2$ *Thermodynamics:* $K_{avg} = \frac{3}{2}k_B T$	*Atomic and Nuclear:* $K_{max} = hf - \phi$
Potential Energy (J)	$\Delta U_g = mgh$ $U_G = -\frac{Gm_1m_2}{r}$ $G = 6.67 \times 10^{-11}\ \frac{Nm^2}{kg^2}$	*Coiled Spring:* $U_s = -\frac{1}{2}k\Delta x^2$	$U_E = qV$ $U_E = \frac{1}{4\pi\epsilon_0}\frac{q_1q_2}{r}$ $U_c = \frac{1}{2}QV$ $U_c = \frac{1}{2}CV^2$ $U_L = \frac{1}{2}LI^2$	*Fluid Mechanics:* $P = P_0 + \rho g h$ *Continuity of Mass:* $\rho_1 A_1 v_1 = \rho_2 A_2 v_2$ *Continuity of Volume:* $A_1 v_1 = A_2 v_2$ Thermodynamics: $\Delta U = Q + W$	*Atomic and Nuclear:* $E = hf$ $E = mc^2$ $\Delta E = (\Delta m)c^2$ *Relativity:* $E = mc^2\left(\frac{1}{\sqrt{1 - \frac{v^2}{c^2}}}\right)$

Copyright © 2011-2023 by Harold Toomey, Wyzant Tutor

	Mechanics: Linear	Mechanics: Angular	Electricity / Magnetism	Fluid Mechanics / Thermodynamics	Atomic and Nuclear / Waves and Optics
Heat Energy (J)	*Conservation of Energy:* $E_i = E_f$ $\sum E_i = \sum E_f$ $E = W + Q + K + \Delta U_g + U_G + U_s + U_E + U_c + U_L + \cdots = constant$			*<u>Thermodynamics</u>:* $H = \frac{kA\Delta T}{L}$ $Q = mH_f$ $Q = mH_v$ $\Delta E = Q = mC\Delta T$ $mC\Delta T = mC(T_f - T_i)$ $C_{H_2O} = 4180\,J\,\frac{J}{kg\,°K}$ $T_f = \frac{m_1C_1\Delta T_{1i} + m_2C_2\Delta T_2}{m_1C_1 + m_2C_2}$	*NA*
Power (W)	$P = \frac{W}{t} = Fv$ $P = \frac{\Delta E}{\Delta t} \rightarrow \frac{dE}{dt}$ $P = Fv\cos\theta$ $P = \boldsymbol{F} \bullet \boldsymbol{v}$	$P = \frac{W}{t} = \tau\omega$ $P = \frac{\Delta W}{\Delta t} \rightarrow \frac{dW}{dt}$ $P = \tau\omega\cos\theta$ $P = \boldsymbol{\tau} \bullet \boldsymbol{\omega}$	$P = IV$ $P = I^2R$ $P = \frac{V^2}{R}$	$P = pQ$ *where:* $P = pressure\left(\frac{N}{m^2}\right)$ $Q = volumetric\ flow\ rate\left(\frac{m^3}{s}\right)$	*NA*

Copyright © 2011-2023 by Harold Toomey, Wyzant Tutor

	Mechanics: Linear	Mechanics: Angular	Electricity / Magnetism	Fluid Mechanics / Thermodynamics	Atomic and Nuclear / Waves and Optics
Engineering Application					
Period / Frequency (Hz)	$T = \frac{1}{f}$ $f = \frac{1}{T}$ *Kepler's Third Law:* $T^2 = \left(\frac{4\pi^2}{GM}\right) r^3$	$T = \frac{1}{f} = \frac{2\pi}{\omega}$ $T_s = 2\pi\sqrt{\frac{m}{k}}$ $T_p = 2\pi\sqrt{\frac{\ell}{g}}$	$T = \frac{2\pi}{b}$ *For:* $y = sin(b\theta)$ $y = cos(b\theta)$	*NA*	*<u>Waves and Optics</u>:* $f = \frac{v}{\lambda}$ *Doppler Effect:* $\boldsymbol{f_r = f_s\left(\frac{v \pm v_r}{v \mp v_s}\right)}$
Center of Mass (m)	$x_{CM} = \frac{m_1 x_1 + m_2 x_2}{m_1 + m_2}$ $x_{CM} = \frac{\sum m_i x_i}{\sum m_i}$ $\bar{x} = \frac{1}{M}\int_0^M x\, dm$ *where* $M = \int_0^M dm$ *and* $dm = \rho\, dz\, dy\, dx$	$\boldsymbol{r}_{CM} = \frac{\sum m\boldsymbol{r}}{\sum m}$	*NA*	*NA*	*NA*
Rigid Bodies	$\sum F_y = \sum mg = 0$ *(Down = '–')*	$\sum \tau = \sum F_y\, x_{CM} = 0$ *(CW = '–')*	*NA*	*NA*	*NA*

Copyright © 2011-2023 by Harold Toomey, Wyzant Tutor

	Mechanics: Linear	Mechanics: Angular	Electricity / Magnetism	Fluid Mechanics / Thermodynamics	Atomic and Nuclear / Waves and Optics
Conservation Laws	*Conservation of Linear Momentum* $\sum p_i = \sum p_f$ $\sum m\boldsymbol{v}_i = \sum m\boldsymbol{v}_f$ *(Physics)*	*Conservation of Angular Momentum* $\sum L_i = \sum L_f$ $\sum I\boldsymbol{\omega}_i = \sum I\boldsymbol{\omega}_f$ *(Physics)*	*Conservation of Electric Charge* $\sum Q_i = \sum Q_f$ *(Circuits)*	*Conservation of Mass (or Matter)* $\sum m_i = \sum m_f$ *(Chemistry)* *Bernoulli's Equation:* $P_1 + \rho g y_1 + \frac{1}{2}\rho v_1^2 = P_2 + \rho g y_2 + \frac{1}{2}\rho v_2^2$ *(Fluid Mechanics / Pressure)*	*Conservation of Energy* $\sum E_i = \sum E_f$ $E = W + Q + K + \Delta U_g + U_G + U_s + U_E + U_c + U_L + \cdots = constant$ *(Physics)*

Copyright © 2011-2023 by Harold Toomey, Wyzant Tutor

Electricity			
Electricity			
Electric Field (V/m) (N/C)	$\boldsymbol{E} = \frac{\boldsymbol{F}}{q}$ $E_{avg} = -\frac{V}{d}$ $E = -\frac{\Delta V}{\Delta r} \rightarrow -\frac{dV}{dr}$ *Gauss's Law:* $\oint \boldsymbol{E} \bullet d\boldsymbol{A} = \frac{Q}{\epsilon_0}$ $\boldsymbol{E} = \rho \boldsymbol{J}$	**Resistance (Ω)**	$R = \frac{V}{I}$ $R = \frac{\rho \ell}{A}$ $R_s = \sum_i R_i$ $\frac{1}{R_p} = \sum_i \frac{1}{R_i}$ $R_p = \frac{R_1 R_2}{R_1 + R_2}$
Potential / Voltage (V)	$V = IR$ $V = \frac{Q}{C}$ $V = k \sum_i \frac{q_i}{r_i}$ $V = \frac{1}{4\pi\epsilon_0} \sum_i \frac{q_i}{r_i}$	**Capacitance (F)**	$C = \frac{Q}{V}$ $C = \frac{\epsilon_0 A}{d}$ $C = \frac{\kappa \epsilon_0 A}{d}$ $C_p = \sum_i C_i$ $\frac{1}{C_s} = \sum_i \frac{1}{C_i}$
Current (A)	$I = \frac{\Delta Q}{\Delta t} \rightarrow \frac{dQ}{dt}$ $I = \frac{V}{R}$ $I = Nev_d A$ $e = -1.602176462 \times 10^{-19}\ C$	**Inductance (H)**	$L = N\frac{\Phi}{I}$ $L = \frac{V_L}{dI/dt}$ $L = \mu_0 \frac{N^2 A}{\ell}$ $L_s = \sum_i L_i$ $\frac{1}{L_p} = \sum_i \frac{1}{L_i}$

Copyright © 2011-2023 by Harold Toomey, Wyzant Tutor

Magnetism			
Magnetism			
Magnetic Field (T)	$B = \frac{\mu_0}{2\pi}\frac{I}{r}$ $Solenoid:$ $B_s = \mu_0 n I$ $where\ n = \frac{N}{\boldsymbol{\ell}}\ turns\ per\ meter$ $\boldsymbol{B} = \frac{\mu_0}{4\pi}\frac{q\vec{\boldsymbol{v}} \times \vec{\boldsymbol{r}}}{r^3}$ $\boldsymbol{B} = \frac{\mu_0}{4\pi}\frac{q\vec{\boldsymbol{v}} \times \hat{\boldsymbol{r}}}{r^2}$ $\mu_0 = 4\pi \times 10^{-7}\frac{(T \bullet m)}{A}$ *Ampere's Circuit Law:* $\oint \boldsymbol{B} \bullet d\boldsymbol{\ell} = \mu_o I + \mu_0 \varepsilon_0 \frac{d\phi_E}{dt}$ *Gauss's Law for Magnetism:* $\oint \boldsymbol{B} \bullet d\boldsymbol{A} = 0$ $dB = \frac{\mu_0}{4\pi}\frac{I\, d\boldsymbol{\ell} \times \vec{\boldsymbol{r}}}{r^3}$	**EMF (V)**	*Faraday's Law of Induction:* $\oint \boldsymbol{E} \bullet d\boldsymbol{\ell} = -\frac{\Delta\phi_B}{\Delta t}$ $\varepsilon = -L\frac{dI}{dt}$
Magnetic Flux (Wb)	$\phi_{\boldsymbol{B}} = BA\cos\theta$ $\phi_B = \boldsymbol{B} \bullet \boldsymbol{A}$ *Gauss's Law for Magnetism:* $\phi_B = \int \boldsymbol{B} \bullet d\boldsymbol{A}$		
EMF (V)	$\varepsilon_{avg} = -\frac{\Delta\phi_B}{\Delta t}$ $\boldsymbol{\varepsilon} = \oint \boldsymbol{E} \bullet d\boldsymbol{s} = -\frac{d\phi_B}{dt}$ $\boldsymbol{\varepsilon} = \boldsymbol{B}lv$		

Copyright © 2011-2023 by Harold Toomey, Wyzant Tutor